

An Euler-Type Formula for $\zeta(2k + 1)$

Michael J. Dancs and Tian-Xiao He

Department of Mathematics and Computer Science

Illinois Wesleyan University

Bloomington, IL 61702-2900, USA

Draft, June 30th, 2004

Abstract

In this short paper, we give several new formulas for $\zeta(n)$ when n is an odd positive integer. The method is based on a recent proof, due to H. Tsumura, of Euler's classical result for even n . Our results illuminate the similarities between the even and odd cases, and may give some insight into why the odd case is much more difficult.

AMS Subject Classification: 11M06, 11Y35, 41A30, 65D15.

Key Words and Phrases: Riemann Zeta function, Euler's formula, Euler polynomial, Bernoulli number.

1 Introduction

Let $\zeta(s)$ be the Riemann zeta function. In [1], Tsumura gave an elementary proof of Euler's well-known formula

$$\zeta(2k) = \frac{(-1)^{k-1} 2^{2k-1} \pi^{2k}}{(2k)!} B_{2k} \quad (1.1)$$

where k is a positive integer and $\{B_n\}$ denotes the sequence of Bernoulli numbers. In this paper, we use Tsumura's method to develop an "Euler-type" formula for $\zeta(2k + 1)$ analogous to (1.1) above.

2 Preliminaries

For $d > 0$ and $u \in [1, 1 + d]$, we let

$$\frac{2e^t}{e^t + u} = \sum_{n=0}^{\infty} \phi_n(u) \frac{t^n}{n!}, \quad (2.1)$$

We observe that $\phi_n(1) = E_n(1)$, where $E_n(x)$ is the n th Euler polynomial. If n is a nonnegative integer and $u > 1$, we have

$$\phi_n(u) = -2 \sum_{j=1}^{\infty} (-u)^{-j} j^n. \quad (2.2)$$

When n is a negative integer, we take (2.2) as our definition of $\phi_n(u)$. It is easily shown that $\phi_{-1}(1) = 2 \ln 2$, and that

$$\phi_{-m}(1) = -2(2^{1-m} - 1)\zeta(m) \quad (2.3)$$

whenever m is an integer greater than 1. Finally, we note that for $u \geq 1$,

$$\overline{\lim}_{n \rightarrow \infty} \left(\frac{|\phi_n(u)|}{n!} \right)^{1/n} \leq \frac{1}{\pi}, \quad (2.4)$$

and thus the series in Eq. (2.1) converges absolutely for $|t| < \pi$.

For any positive integer k , we have

$$\begin{aligned} 0 &= \sum_{n=1}^{\infty} \frac{(-u)^{-n} \sin(n\pi)}{n^{2k}} \\ &= \sum_{n=1}^{\infty} \frac{(-u)^{-n}}{n^{2k}} \sum_{j=0}^{\infty} (-1)^j \frac{(n\pi)^{2j+1}}{(2j+1)!} \\ &= \sum_{j=0}^{\infty} \frac{(-1)^{j+1} u \pi^{2j+1}}{2(2j+1)!} \phi_{2j+1-2k}(u) \\ &= \sum_{j=0}^{k-1} \frac{(-1)^{j+1} u \pi^{2j+1}}{2(2j+1)!} \phi_{2j+1-2k}(u) + \sum_{j=k}^{\infty} \frac{(-1)^{j+1} u \pi^{2j+1}}{2(2j+1)!} \phi_{2j+1-2k}(u) \\ &= \sum_{j=0}^{k-1} \frac{(-1)^{j+1} u \pi^{2j+1}}{2(2j+1)!} \phi_{2j+1-2k}(u) + \sum_{m=0}^{\infty} \frac{(-1)^{m+k+1} u \pi^{2m+2k+1}}{2(2m+2k+1)!} \phi_{2m+1}(u). \end{aligned}$$

In light of (2.4), we can now let $u \rightarrow 1^+$, obtaining

$$0 = \sum_{j=0}^{k-1} \frac{(-1)^{j+1} \pi^{2j+1}}{2(2j+1)!} \phi_{2j+1-2k}(1) + \sum_{m=0}^{\infty} \frac{(-1)^{k+1} \pi^{2k+1} f_m}{2(2m+2k+1)!}, \quad (2.5)$$

where $f_m = (-1)^m \pi^{2m} E_{2m+1}(1)$.

Setting $k = 1$ in (2.5) and recalling that $\phi_{-1}(1) = 2 \ln 2$, we have the curious formula

$$\ln 2 = \frac{\pi^2}{2} \sum_{m=0}^{\infty} \frac{(-1)^m \pi^{2m}}{(2m+3)!} E_{2m+1}(1). \quad (2.6)$$

3 Main Results

We can use (2.3) and (2.5) to deduce the following theorem, which gives $\zeta(2k+1)$ recursively in terms of $\ln 2$, $\zeta(3), \dots, \zeta(2k-1)$:

Theorem 3.1 *For any positive integer k ,*

$$\begin{aligned} (1 - 2^{-2k}) \zeta(2k+1) &= \sum_{j=1}^{k-1} \frac{(-1)^j \pi^{2j}}{(2j+1)!} (2^{2j-2k} - 1) \zeta(2k-2j+1) \\ &\quad - \frac{(-1)^k \pi^{2k} \ln 2}{(2k+1)!} + \frac{(-1)^k \pi^{2k+2}}{2} \sum_{m=0}^{\infty} \frac{(-1)^m \pi^{2m} E_{2m+1}(1)}{(2m+2k+3)!}. \end{aligned} \quad (3.1)$$

This result may be regarded as the analogue of equation (5) of [1], in the sense that the infinite series above reduces to a single term if the E_{2m+1} is replaced by an E_{2m} . This provides some perspective on the difficulty of evaluating $\zeta(2k+1)$ as opposed to $\zeta(2k)$.

When $k = 1, 2$, and 3 , Theorem 3.1 yields the respective formulas:

$$\begin{aligned} \zeta(3) &= \frac{\pi^2}{9} \ln 4 - \frac{2\pi^4}{3} \sum_{m=0}^{\infty} \frac{(-1)^m \pi^{2m} E_{2m+1}(1)}{(2m+5)!}, \\ \zeta(5) &= \frac{2\pi^2}{15} \zeta(3) - \frac{\pi^4}{225} \ln 4 + \frac{8\pi^6}{15} \sum_{m=0}^{\infty} \frac{(-1)^m \pi^{2m} E_{2m+1}(1)}{(2m+7)!}, \\ \zeta(7) &= \frac{10\pi^2}{63} \zeta(5) - \frac{2\pi^4}{315} \zeta(3) + \frac{\pi^6}{19845} \ln 16 - \frac{32\pi^8}{63} \sum_{m=0}^{\infty} \frac{(-1)^m \pi^{2m} E_{2m+1}(1)}{(2m+9)!}. \end{aligned}$$

If desired, we may express the infinite series in these formulas in terms of the Bernoulli numbers, via the identities

$$E_{2m+1}(1) = -E_{2m+1}(0) = \frac{2(2^{2m+2} - 1)}{2m + 2} B_{2m+2}.$$

It is well-known that

$$|B_{2m+2}| < \frac{2(2m + 2)!}{(2\pi)^{2m+2} (1 - 2^{-2m-1})},$$

and this implies that the m th term of the series in (3.1) is $O(m^{-2k-2})$. Hence our result gives slightly faster convergence than the standard series for $\zeta(2k + 1)$.

Building on Theorem 3.1, we now develop the following Euler-type formula for $\zeta(2k + 1)$:

Theorem 3.2 *Let $f_m = \pi^{2m}(-1)^m E_{2m+1}(1)$. For any positive integer k ,*

$$\zeta(2k + 1) = \frac{(-1)^{k+1} \pi^{2k+2}}{(1 - 2^{-2k})} \sum_{m=0}^{\infty} \frac{P_k(m) f_m}{(2m + 2k + 3)!}, \quad (3.2)$$

where $P_k(m)$ is a polynomial in m with rational coefficients, having degree at most $2k$. For $k \geq 0$, we have the recurrence:

$$(-1)^{k+1} P_k(m) = \frac{1}{2(m + k + 2)} \sum_{l=0}^{k-1} (-1)^l \binom{2m + 2k + 4}{2m + 2l + 3} P_l(m) - \frac{1}{2}. \quad (3.3)$$

Proof. For ease of notation, we set $Z(s) = \frac{1}{2} \phi_{-(2s+1)}(1)$, noting that $Z(0) = \ln 2$ and $Z(n) = (1 - 2^{-2n}) \zeta(2n + 1)$ for any positive integer n . We may rewrite (3.1) as

$$Z(k) = - \sum_{j=1}^k \frac{(-1)^j \pi^{2j}}{(2j + 1)!} Z(k - j) + \frac{(-1)^k \pi^{2k+2}}{2} \sum_{m=0}^{\infty} \frac{f_m}{(2m + 2k + 3)!}. \quad (3.4)$$

If $Z(k) = \pi^{2k+2} \sum_{m=0}^{\infty} P_k(m) f_m / ((2m + 2k + 3)!)$, we see from (2.6) that $P_0(m) = 1/2$. For $k > 0$ we have

$$\begin{aligned}
Z(k) &= -\sum_{j=1}^k \frac{(-1)^j \pi^{2j}}{(2j+1)!} \pi^{2k-2j+2} \sum_{m=0}^{\infty} \frac{P_{k-j}(m) f_m}{(2m+2k-2j+3)!} \\
&\quad + \frac{(-1)^k \pi^{2k+2}}{2} \sum_{m=0}^{\infty} \frac{f_m}{(2m+2k+3)!} \\
&= -\pi^{2k+2} \sum_{j=1}^k \frac{(-1)^j}{(2j+1)!} \sum_{m=0}^{\infty} \frac{P_{k-j}(m) f_m}{(2m+2k-2j+3)!} \\
&\quad + \frac{(-1)^k}{2} \sum_{m=0}^{\infty} \frac{f_m}{(2m+2k+3)!} \\
&= -\pi^{2k+2} \sum_{l=0}^{k-1} \frac{(-1)^{k-l}}{(2k-2l+1)!} \sum_{m=0}^{\infty} \frac{P_l(m) f_m}{(2m+2l+3)!} + \sum_{m=0}^{\infty} \frac{f_m}{2(2m+2k+3)!} \\
&= (-1)^{k+1} \pi^{2k+2} \sum_{m=0}^{\infty} \left(\sum_{l=0}^{k-1} \frac{(-1)^l P_l(m) f_m}{(2k-2l+1)!(2m+2l+3)!} - \frac{f_m}{2(2m+2k+3)!} \right).
\end{aligned}$$

Hence

$$\begin{aligned}
(-1)^k P_k(m) &= \frac{1}{2} - \sum_{l=0}^{k-1} \frac{(-1)^l P_l(m) (2m+2k+3)!}{(2k-2l+1)!(2m+2l+3)!} \\
&= \frac{1}{2} - \frac{1}{2(m+k+2)} \sum_{l=0}^{k-1} (-1)^l P_l(m) \binom{2m+2k+4}{2m+2l+3}. \quad (3.5)
\end{aligned}$$

■

Equation (3.3) can be written in the following closed form as a partition of unity:

$$\frac{1}{m+k+2} \sum_{\ell=0}^k (-1)^\ell P_\ell(m) \binom{2m+2k+4}{2m+2\ell+3} = 1.$$

The first few $P_k(m)$ are given by

$$\begin{aligned}
P_1(m) &= \frac{(m+1)(2m+7)}{6}, \\
P_2(m) &= \frac{(m+1)(m+2)(28m^2+224m+465)}{180}, \\
P_3(m) &= \frac{(m+1)(m+2)(m+3)(248m^3+3348m^2+15346m+24003)}{3780},
\end{aligned}$$

and we can use these to obtain formulas for $\zeta(3)$, $\zeta(5)$ and $\zeta(7)$, respectively. The pattern suggested by the above formulas holds in general, and we have:

Theorem 3.3 *Let $P_k(m)$ be defined as in Theorem 3.2. For any integers k and n with $1 \leq n \leq k$, $P_k(-n) = 0$.*

Proof. We first establish

$$P_{n-1}(-n) = \frac{(-1)^{n-1}}{2} \quad (3.6)$$

for any positive integer n . From (3.3) we have

$$(-1)^{n+1}P_n(-n-1) = \frac{1}{2} \sum_{\ell=0}^{n-1} (-1)^\ell \binom{2}{2\ell-2n+1} P_\ell(-n-1) - \frac{1}{2} = -\frac{1}{2},$$

which implies (3.6).

We now proceed by induction on k . Assuming that $P_\ell(-n) = 0$ when $1 \leq n \leq \ell \leq k$, we must show that $P_{k+1}(-n) = 0$ when $1 \leq n \leq k+1$. From (3.3) we have

$$\begin{aligned} (-1)^{k+2}P_{k+1}(-n) &= \frac{1}{2(k-n+3)} \sum_{\ell=0}^k (-1)^\ell \binom{2k-2n+6}{2\ell-2n+3} P_\ell(-n) - \frac{1}{2} \\ &= \frac{1}{2(k-n+3)} \sum_{\ell=n-1}^k (-1)^\ell \binom{2k-2n+6}{2\ell-2n+3} P_\ell(-n) - \frac{1}{2}. \end{aligned}$$

Since $P_\ell(-n) = 0$ if $n \leq \ell \leq k$, the right-hand side is

$$\frac{1}{2(k-n+3)} (-1)^{n-1} (2k-2n+6) P_{n-1}(-n) - \frac{1}{2} = 0. \quad (3.7)$$

■

In light of Theorem 3.3 and computational evidence, we propose the following conjecture.

Conjecture 1 *For any positive integer k , the polynomial $P_k(m)$ has simple roots $m = -1, -2, \dots, -k$, and no other rational roots if $k \geq 2$.*

Although the Euler-type formulas from Theorem 3.2 are more compact, they converge very slowly as compared to Theorem 3.1. As a compromise, we give:

Theorem 3.4 *Let $f_m = (-1)^m \pi^{2m} E_{2m+1}(1)$. For any positive integer k ,*

$$\zeta(2k+1) = \frac{(-1)^k \pi^{2k}}{1-2^{-2k}} \left[a_k \ln 2 - \pi^2 \sum_{m=0}^{\infty} \frac{Q_k(m) f_m}{(2m+2k+3)!} \right], \quad (3.8)$$

where a_k is a constant and $Q_k(m)$ is a polynomial in m with rational coefficients, having degree at most $2k-2$. Recursive formulas for a_k and $Q_k(m)$ are given by

$$a_k = - \sum_{\ell=1}^{k-1} \frac{a_{\ell}}{(2k-2\ell+1)!} - \frac{1}{(2k+1)!} \quad (3.9)$$

and

$$Q_k(m) = - \frac{1}{2(m+k+2)} \sum_{\ell=1}^{k-1} \binom{2m+2k+4}{2m+2\ell+3} Q_{\ell}(m) - \frac{1}{2}. \quad (3.10)$$

The proof of this result is similar to that of Theorem 3.2, and hence is omitted.

Eqs. (3.9) and (3.10) can be written in closed form as

$$-(2k+1)! \sum_{\ell=1}^k \frac{a_{\ell}}{(2k-2\ell+1)!} = 1$$

and

$$- \frac{1}{m+k+2} \sum_{\ell=1}^k Q_{\ell}(m) \binom{2m+2k+4}{2m+2\ell+3} = 1,$$

respectively. For $k=1, 2$, and 3 , we obtain $a_1 = -1/6$, $a_2 = 7/360$, $a_3 = -31/15120$, $Q_1(m) = -1/2$, $Q_2(m) = (m+2)(2m+9)/6$, and $Q_3(m) = -(m+2)(m+3)(28m^2+280m+717)/180$. Hence,

$$\begin{aligned} \zeta(3) &= \frac{\pi^2}{9} \ln 4 - \frac{2\pi^4}{3} \sum_{m=0}^{\infty} \frac{f_m}{(2m+5)!}, \\ \zeta(5) &= \frac{7\pi^4}{675} \ln 4 - \frac{8\pi^6}{45} \sum_{m=0}^{\infty} \frac{(m+2)(2m+9)}{(2m+7)!} f_m, \\ \zeta(7) &= \frac{62\pi^6}{59535} \ln 4 - \frac{16\pi^8}{2835} \sum_{m=0}^{\infty} \frac{(m+2)(m+3)(28m^2+280m+717)}{(2m+9)!} f_m. \end{aligned}$$

Additionally, we have the following result on the roots of $Q_k(m)$:

Theorem 3.5 *Let k and n be integers. Then the polynomials $Q_k(m)$ satisfy $Q_n(-n-1) = -\frac{1}{2}$ for all $n \geq 1$ and $Q_k(-n) = 0$ for all $k \geq n \geq 2$.*

We may also make the following:

Conjecture 2 *For $k \geq 2$, the polynomial $Q_k(m)$ has simple roots $-2, -3, \dots, -k$, and $Q_k(m)$ has no other rational roots if $k \geq 3$.*

References

- [1] H. Tsumura, An elementary proof of Euler's formula for $\zeta(2m)$, *American Mathematical Monthly*, **111**(2004), 430-431.